

Clinical References for Fibro Eze

1. Abraham GE, Flechas JD. Hypothesis: Management of fibromyalgia: rational for the use of magnesium and malic acid. *J Nutr Med* 3:49-59, 1991
2. Caruso I, et al. Double-blind study of 5-hydroxytryptophan versus placebo in the treatment of primary fibromyalgia syndrome. *J Int Med Res* 18 (3): 201-209.
3. Eisinger J, et al. Donnees actuelles sur les fibromyalgies: traitement des fibromyalgies primitives par la cocarboxylase. *Lyon Med Med* 24: 11585-6, 1988 (in French)
4. Eisinger J, et al. Glycolysis abnormalities in fibromyalgia *J Am Coll Nutr* 13 (2): 144-148, 1994
5. Eisinger J., et al. Selenium and magnesium status in fibromyalgia. *Magnes Res* 7(3-4):285-288, 1994.
6. Moldofsky H, Warsh JJ. Plasma tryptophan and musculoskeletal pain in non-articular rheumatism. *Pain* 5(1):65-71, 1978.
7. Romano TJ, Stiller JW. Magnesium deficiency in fibromyalgia syndrome. *J Nutr Med* 4:165-167, 1994.
8. Yunus B, et al. Plasma tryptophan and other amino acids in primary fibromyalgia: acontrolled study. *J Rheumatol* 19(1):90-94, 1992.